


Checklist of Hypogonadism Symptoms and Comorbidities

The following symptoms and comorbidities are associated with hypogonadism. Based on the individual patient's history, further testing may be needed when several of the following present in combination.

SYMPTOMS



COMORBIDITIES

- Anemia
- CVD
- Obesity
- Metabolic syndrome, type 2 diabetes
- Osteoporosis, low BMD
- Dyslipidemia
- Hypertension
- HIV-associated weight loss
- End-stage renal disease and maintenance hemodialysis
- Moderate to severe COPD

BMD, bone mineral density; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CVD, cardiovascular disease; ED, erectile dysfunction; HIV, human immunodeficiency virus.

Bibliography

- Bhasin S, Cunningham GR, Hayes FJ, et al. Testosterone therapy in adult men with androgen deficiency syndromes: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* 2006;91(6):1995-2010.
- Guay A, Seftel AD, Traish A. Hypogonadism in men with erectile dysfunction may be related to a host of chronic illnesses. *Int J Impot Res.* 2010;22(1):9-19.
- Sadovsky R. It's time for a large, long-term study of testosterone supplementation in hypogonadal older men: a primary care view. *J Mens Health Genet.* 2006;3(1):36-38.
- Seftel A. Male hypogonadism. Part II: etiology, pathophysiology, and diagnosis. *Int J Impot Res.* 2006;18(3):223-228.

Jointly sponsored by the Annenberg Center for Health Sciences at Eisenhower and CogniMed Inc. This information sheet is supported by an independent educational grant provided by Endo Pharmaceuticals.